RedoxAG



# **Guaranteed Analysis**

Boron (B) . . . . . . . . . . . . 2.0%\* Copper (Cu) . . . . . . . . . . 2.0% Iron (Fe) . . . . . . . . . . . 6.0% Manganese (Mn) . . . . . . . . 6.0% 6%.....Water Soluble Manganese Molybdenum (Mo) . . . . . . . 0.001%\*\*

Zinc (Zn). . . . . . . . . . . . 6.0% (Derived from Boric Acid, Copper Sulfate, Ferric Sulfate, Manganese Sulfate, Sodium Molybdate and Zinc Sulfate)

F001388 \*WARNING Boron: The use of the fertilizing material on any crops other than those recommended may result in serious injury to the crop(s)
\*\* WARNING Molybdenum: The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

## **KEEP OUT OF REACH OF CHILDREN**

### STOP, READ THE LABEL BEFORE USING THIS PRODUCT

WARRANTY: Since weather, crops, soil, and other conditions may vary, the seller makes no warranty of any kind, express or implied, concerning this product. The user assumes all risk of use and handling whether or not in accordance with directions or suggestions.

### **COMPATIBILITY**

Always check for compatibility with other products prior to use.

#### **GENERAL INFORMATION**

TriPlex Micro is a dry soluble product, developed as an easy to use formulation that is compatible with most fertilizers and foliar sprays.

#### **APPLICATION RATES**

Application	Rate	Comments
Foliar Spray	1-3 lbs/acre Or 0.4-1.2 oz/1000 ft <sup>2</sup>	Maintain a 1% solution by weight or less if material might come in contact with highly sensitive plants. Includes all other TriPlex products     Do not exceed 2 lbs/acre if applied in less than 25/gal/acres
Soil	1-6 lbs/acre Or 0.4-2.2 oz 1000 ft <sup>2</sup>	> Apply with a minimum of 15 gallons of water/acre

### APPLICATION GUIDELINES

- Use extreme caution when applying foliar nutrients to fruit that may be susceptible to marking. Avoid using Redox TriPlex foliar nutrient fertilizers with penetrating surfactants on fruit that may be susceptible to marking.

  Proper application rates and frequency should be determined by specific conditions and plant requirements.

  Apply during periods of high micronutrient demand.

  Always jar test before adding to a spray tank or injection tank.

  Spray applications should be made with sufficient water, 50 mesh screens, and agitation throughout the application.

  Do not spray on highly sensitive plant material at a rate of more than 1% by weight.

  When tank mixing with other chamicals, it is the responsibility of the and user to assume compatibility and safety.

- When tank mixing with other chemicals, it is the responsibility of the end-user to assume compatibility and safety.
- Poly in-line filters should be replaced with stainless steel, if possible.
- Always consult your agronomist.

AVOID CONTACT WITH SKIN, EYES OR CLOTHING

In case of contact immediately flush skin, eyes or clothing with plenty of water for several minutes. Distributed and Guaranteed by: Manufactured by: Redox Chemicals, LLC



QUIMICA CREATIVA A SU SERVICIO COSMOCEL, S.A.

Vía a Matamoros No. 1501, San Nicolás de los Garza, N.L., México C.P. 66484 Tel. +52 (81) 8625-3100 www.cosmocel.com servicioaclientes@cosmocel.com



NET CONTENTS: 20 Lbs. (9.08 Kg)



130 S. 100 W. Burley, Idaho, USA 83318 208-678-2610 www.redoxchem.com

08/14

Caution: The products manufactured by Redox Chemicals, LLC are sold only thru licensed distributors. To insure proper mixing and application, consult with trained Redox Chemicals, LLC distributors. Visit www.redoxchem.com for a list of trained and authorized Redox Chemicals, LLC distributors.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm